



Colonic Plasmacytoma Revealing Kahler's Disease: A Case Report

Armandaind M. H. Yila^{1*}, Mohamed Amine El Mansouri¹, Soumaya Jamil¹, Bahya Jarni¹, Wafae Ait Belaid¹, Tariq Ahbala¹, Khalid Rabbani¹, Abdelouahed Louzi¹

¹Department of Visceral Surgery, Mohammed VI University Hospital, Faculty of Medicine and Pharmacy, Cadi Ayyad University, Marrakech, Morocco

*Corresponding author: Armandaind M. H. Yila

Department of Visceral Surgery, Mohammed VI University Hospital, Faculty of Medicine and Pharmacy, Cadi Ayyad University, Marrakech, Morocco

Article History

Received: 30-05-2025

Accepted: 09-05-2025

Published: 16-05-2025



Summary:

Plasmacytoma is defined as a malignant differentiation of B lymphocytes with production of serum immunoglobulins. Plasmacytoma can affect most organs either as a secondary location of a multiple myeloma or as a primary location called extramedullary plasmacytoma. We report a rare case of colonic plasmacytoma revealing Kahler's disease.

Keywords: Colonic plasmacytoma, kahler's disease, intestinal occlusion.

Case Report

Copyright © 2025 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

Plasmacytoma is a monoclonal proliferation of malignant plasma cells that may develop in isolation, corresponding to solitary medullary or extramedullary plasmacytoma, or as part of multiple myeloma [1].

Secondary sites of Kahler's disease most commonly involve the respiratory tract in around 75% of cases. In contrast, the gastrointestinal tract is involved in only 5-10% of cases [2], with colonic involvement being extremely rare [3]. We report a rare case of colonic plasmacytoma revealing Kahler disease.

OBSERVATION

A 65-year-old patient with no specific pathological history presented to the emergency department with abdominal pain in the IDF and right flank, associated with cessation of bowel movements and gas and vomiting, which had been progressing for 5 days. The patient also had asthenia and anorexia associated with significant weight loss. On clinical examination, the patient was in poor general condition; the abdomen was distended and tympanic; the lymph nodes were free; on rectal examination, the rectal ampulla was empty; the rest of the clinical examination was unremarkable.

The ASP carried out had objectified NHA with a predominance of bowel, the patient had then benefited from an abdominal CT scan demonstrating distension of the bowel upstream of a circumferential and

irregular thickening 33mm thick at the level of the ileo-caecal junction associated with some mesenteric adenopathy at the level of the IDF (Fig 1).

The patient benefited from a carcinological right hemicolectomy with ileo-

colic anastomosis. Histological examination of the resection specimen objective plasmacytic infiltration of less than 10%, with infiltration by large blast cells with an appearance of plasmoblasts or immunoblasts (Fig 2).




Figure 1: Abdominal-pelvic CT image showing hydroaeric distension of the ileal ansae upstream of irregular thickening of the ileo-caecal junction




Figure 2: Hemalun-eosin colonic biopsy, magnification x 10 Dense focal plasma cell infiltration of the chorion dissociating the gland (left) and x 40 immature plasmoblastic or immunoblastic cells & the bone marrow biopsy

DISCUSSION

The digestive sites of Kahler disease [3] involve, in decreasing order of frequency, the stomach, jejunum, ileum, colon, rectum, duodenum and oesophagus. Patients may present with several digestive tract sites [4]. However, colonic involvement is exceptional.

Intestinal and colorectal involvement usually leads to symptoms of occlusion, as in the case of our patient, or digestive haemorrhage. Intestinal invagination and

ileocolic fistula have also been described [5, 6]. However, colonic localisation may be asymptomatic.

Digestive plasmacytomas are most often polyploid and appear as adenocarcinomas. Endoscopic polypectomy followed by radiotherapy appears to be the logical treatment. The case report presents an ulcerative form of colonic plasmacytoma not suitable for endoscopic polypectomy.

Colonoscopy has a great place in reducing morbidity, mortality and the cost of medical care when endoscopic polypectomy is possible without laparotomy.

All lesions apart from those that are sessile and large can be completely resected endoscopically [3]. In general, colonic plasmacytoma progresses slowly, but there is a strong tendency for local relapse [2, 7].

Surgical excision supplemented by radiotherapy has been the recommended treatment for localised plasmacytoma [3, 8, 9].

CONCLUSION

The colonic plasmacytoma of Kahler disease is exceptional; it is most often polyploid and symptomatic. However, it can also be asymptomatic, in which case colonoscopy, in conjunction with biopsies, is of vital importance in making the diagnosis.

There is no established treatment protocol: radiotherapy alone or major surgery combined with radiotherapy and/or chemotherapy, or endoscopic polypectomy supplemented by adjuvant treatment. Only multicentre, comparative studies and prolonged follow-up will be able to determine the advantages of each of these therapeutic approaches.

REFERENCES

1. Brown, C. R., & Liber, A. F. (1939). Multiple plasmoma of the ileum and colon. *Arch Pathol*, 28, 112-3.
2. Wiltshaw, E. (1976). The natural history of extramedullary plasmacytoma and its relation to solitary myeloma of bone and myelomatosis. *Medicine*, 55(3), 217-238.
3. Pais, J. R., Garcia-Segovia, J., Rodriguez-Garcia, J. L., Alvarez-Baleriola, I., & Garcia-Gonzalez, M. (1994). Solitary plasmacytoma of the rectum: report of a case treated by endoscopic polypectomy and radiotherapy. *European journal of surgical oncology: the journal of the European Society of Surgical Oncology*

and the British Association of Surgical Oncology, 20(5), 592-594.

4. Sharma, K. D., & Shrivastav, J. D. (1961). Extramedullary plasmacytoma of gastrointestinal tract-with a case report of plasmoma of rectum and a review of literature. *Archives of Pathology*, 71(1), 229-233.
5. Hampton, J. M., & Gandy, J. R. (1957). Plasmacytoma of the gastro-intestinal tract. *Annals of Surgery*, 145(3), 415-422.
6. Brown, C. R., & Liber, A. F. (1939). Multiple plasmoma of the ileum and colon. *Arch Pathol*, 28, 112-3.
7. Price, A., Quilty, P. M., & Ludgate, S. M. (1987). Extramedullary plasmacytoma of the rectum: two cases treated by radiotherapy. *Clinical radiology*, 38(3), 283-285.
8. Brinch, L., Hannisdal, E., Abrahamsen, A. F., Kvatøy, S., & Langholm, R. (1990). Extramedullary plasmacytomas and solitary plasma cell tumours of bone. *European journal of haematology*, 44(2), 132-135.
9. Frassica, D. A., Frassica, F. J., Schray, M. F., Sim, F. H., & Kyle, R. A. (1989). Solitary plasmacytoma of bone: Mayo Clinic experience. *International Journal of Radiation Oncology* Biology* Physics*, 16(1), 43-48.