

## Marjolin's Ulcer: As an Aggressive Tumor: A Case Report

Dr. Y. Lamaalla<sup>1\*</sup>, Dr. I. Loubna<sup>1</sup>, Dr. Z. Alami<sup>1</sup>, Dr. I. Zine-Eddine<sup>1</sup>, Pr. O. El Atiqi<sup>1</sup>, Dr. S. Azzouzi<sup>1</sup>, Dr. Sylla<sup>1</sup>, Dr. Oudghiri<sup>1</sup>, Pr. Laamrani<sup>1</sup>, Pr. Y. Benchamkha<sup>1,2</sup>

<sup>1</sup>Department of Plastic and Reconstructive Surgery, Mohammed VI University Hospital, Marrakech, Morocco

<sup>2</sup>Faculty of Medicine and Pharmacy, Cadi Ayyad University, Marrakech, Morocco

\*Corresponding author: Dr. Y. Lamaalla

Department of Plastic and Reconstructive Surgery, Mohammed VI University Hospital, Marrakech, Morocco

### Article History

Received: 17-05-2025

Accepted: 24-05-2025

Published: 01-06-2025



### Abstract:

The severity of tissue damage caused by high-voltage electrical injuries to the limbs often complicates the management of such burns. Repair procedures are sometimes ineffective, and amputations become unavoidable. We report the case of a young patient who sustained high-voltage electrical injuries to both upper limbs, resulting in bilateral shoulder disarticulation.

**Keywords:** Electrical injury, high voltage, disarticulation, shoulders.

### Review Article

Copyright © 2025 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

### INTRODUCTION

Marjolin's ulcer (MU) is a rare cutaneous malignancy first described by Dr. Jean-Nicolas Marjolin in 1828 [1]. This entity is well described in the literature as a squamous cell carcinoma (SCC) that develops within a pre-existent cutaneous scar or chronic non-healing wound, such as burn scars that represents most cases, less common settings have included stasis, pilonidal sinuses, chronic venous stasis ulcers, vaccination sites, pressure sores, acne conglobata, osteomyelitis, hidradenitis suppurativa, frost bite, chronic fistulas [2, 3]. MU may develop at any age but tends to affect older individuals.

This is likely due to the typically long latency period from inciting event to malignant transformation, which have ranges 20 to 35 years. A male predilection exists. Lesions occur at any anatomic site, but the lower extremities and the head and neck region are most frequently affected. The

Predominant malignancy in MU is squamous cell carcinoma followed by cases of basal cell carcinoma and malignant melanoma which explains that recently, Marjolin ulcer definition is used synonymous with squamous cell carcinomas (SCC) detected on scar tissues.

The treatment of choice is wide surgical resection. MU is typically an aggressive malignancy with a high recurrence rate and poor 5-year survival. The tumors behave aggressively and have a propensity for local recurrence and lymphnode metastases. Marjolin's ulcers have a high tendency to metastasize [4]. Early recognition and proper staging offers the best chance for cure [5]. There are no confirmed effective protocols for treatment of this disease. We report an uncommon case of an advanced and recurring marjolin's ulcer of the left forearm and wrist of an 47 year-old man.

An 53-year-old Mediterranean man, of low socio- economic level, single, mason by profession, right- handed, having as antecedent a thermal burn by gasoline flame involving the left upper limb specifically the forearm the wrist and part of the hand 52 years ago. On admission, the patient was hemodynamically and respiratorily stable, afebrile, with a normal colored conjunctiva, weight 64 kg without any notion of weight loss. Locally, he presented an ulcerating lesion

on the outer surface of the lower extremity of the forearm and wrist measuring 8cm in long axis, painful, fixed in relation to the superficial and deep planes, bleeding on contact without clinical or radiological adenopathy. An anatomopathological study was requested and came back in favor of a squamous cell carcinoma, considering the patient's history, we suspect that it is a marjolin ulcer on a 52 year old burn.



**Figure 1: Patient admission photos**

Due to the intimate contact of the tumor with the bones of the wrist and the bones of the forearm, and the histological nature of the tumor, and following a multidisciplinary consultation, the decision was to perform a mid-left forearm amputation associated with a homolateral axillary curage. The anatomopathological study came back in favor of a moderately differentiated and infiltrating squamous cell carcinoma measuring 9cm and the resection limit passes in healthy tissue and remains at 11cm, it arrives in contact with the bone but without infiltrating it with a negative curage product 0+/24N.



**Figure 2: Per-operative pictures showing a mid left forearm amputation with homolateral axillary curage was indicated**


The patient was referred to plastic surgery and oncology for regular follow-up, but was lost to follow-up after 4 months. One year after his first operation, the patient presented with an ulcerative lesion on the anterointernal aspect of the left arm, bleeding on contact, painful, measuring 5 cm in length, fixed in relation to the profound plane, evolving for three months according to the patient, thus motivating him to return to our

training.

ABTAP angios can was performed, showing an image of:

- Soft tissue formation of the left scapular region, well limited, oval in shape, with fine partitions and parietal calcifications measuring 55x40 mm, in front: it comes into intimate contact with the deltoid and intraspinous muscles opposite with loss of the separating fatty border, in the back: it comes into contact with the skin covering opposite, without any noticeable skin thickening.
- Left mediastinal and axillary adenopathies of secondary appearance.

- Tissue formations in the posterior soft tissues of the left scapular and dorsal region without any sign of regional invasion. To be compared with histopathological data the decision of the staff between the plastic surgeons, the oncologists and the anaesthetists was to admit the patient to the operating room for a disarticulation of the left limb initially amputated at the level of the forearm with a biopsy of the scapular lesions which turned out to be dermal cysts during the operation. Subcutaneous nodules and necrotic homolateral axillary and supra-clavicular adenopathies suggestive of tumor recurrence.



**Fig 3: A: Recurrence in the anterointernal aspect of the arm 1 year later with two adenopathies, and a pulmonary metastasis made of nodules. B: A Clean-up disarticulation performed followed by adjuvant treatment with chemotherapy was indicated in this patient**

The anatomopathological study of the surgical specimen came back in favor of a moderately differentiated squamous cell carcinoma and infiltrate, the bone is not infiltrated by the tumor, the surgical limits of the soft parts pass in healthy tissue and for the scapular and deltoid cysts it is a reworked fibro-muscular tissue. No signs of malignancy.

The patient was sent directly to oncology for adjuvant treatment with chemotherapy. After two sessions and in half a year, the patient presented a subcutaneous mass in the warm left axilla, fixed in relation to the deep and superficial planes.



**Fig 4: Recurrence of the patient in the left axilla**

An MRI was performed, showing a voluminous locally infiltrating left axillary cystic mass with vascular envelopment associated with Given the non-operability of the lesion and the metastatic nature of the tumor, the patient is a candidate for chemotherapy as part of a palliative treatment.

## DISCUSSION

MU refers to the group of malignant tumors that develop on several types of scars and chronic ulcerations. In most cases, it is a burn scar, as was the case in our patient. In a review of the literature by Mahlon *et al.*, [6], which included 443 patients, 76.5% of these patients had developed MU on old burn scars. MU can also develop on vaccination scars [7], osteomyelitis scars [8], chronic venous stasis scars [9] or pressure sores [10]. Although this malignant transformation of scars has been noted since 100 BC by Celsus, it was not until 1823 that François- Nicolas Marjolin made the first clinical description [11-13]. In most cases, it is a cutaneous squamous cell carcinoma [13-15] as reported in our case. MU usually develops on the limbs and trunk [13] as described in our patient, but also and more particularly on the large flexion folds since the ulceration is sustained by antagonistic muscle forces and frequent trauma. It can also occur in the neck [16] and scalp. The time to onset of these skin tumors varies on average from six weeks [17] to 30 years [18]. The latency period in our patient was 52 years, which is much longer than the figures described in the literature. The etiopathogenesis of DFU is multifactorial and remains poorly elucidated. Indeed, these sequelae of fibrous burns are poorly vascularized, they ulcerate continuously and are very sensitive to micro trauma. Some authors even report the concomitant role of certain mutations, in particular of the Fas gene [15], and of certain local factors such as certain toxins [19]. Castillo and Goldsmith [19] have also shown that immune deficiency could play an important role in this etiopathogeny. On the other hand, most authors agree that it is rarer to encounter neoplasia on operated and graft-covered burn sequelae. Chlihi *et al.*, [11] reported on a series of 54 cases of burn scar development collected over an eight-year period in the burn and plastic surgery department of the Ibn Rochd University Hospital in Casablanca, Morocco, and noted that 78% of the patients had no initial management of the burn. The prognosis of DMU remains guarded given its high metastatic potential, risk of

recurrence, and high mortality and comorbidity [20, 21]. Although the management of DMU is controversial, radical surgery remains the treatment of choice, combined in some cases with radiotherapy and chemotherapy. There is currently no consensus on lymphatic dissection. Most authors agree that prophylactic lymph node dissection is not necessary [22, 23], while Novick *et al.*, [23] emphasize the importance of prophylactic lymph node dissection, especially in cases of lower and upper limb MU. In our case, lymph node dissection was performed, even though the clinical examination and radiological explorations initially showed no lymph node involvement, confirmed by a negative dissection product, only at the recurrence that homo lateral axillary adenopathies were found, hence the interest in discussing the sentinel node and its effectiveness in this type of case. Eastman *et al.*, described successful preoperative lymphoscintigraphy and intraoperative lymphatic mapping in 5 out of their 6 reported patients [24]. None of these patients had palpable lymphadenopathy. Of the 5 patients who underwent successful SLNB, microscopic metastases were identified in 4 of them (80%). The authors note that the prognostic significance remains unclear still, though earlier identification of nodal disease allows for accurate staging and earlier management. In contrast, Motamedolshariati *et al.*, reported only a 2% success rate of sentinel node mapping in their cohort of 10 patients [25]. They attribute this failure to scant lymphatic vessels in the traumatized skin surrounding the Marjolin's Ulcer. Further studies are required to elucidate the indications and rate of success for SLNB. Shen *et al.*, (2014) used preoperative PET-CT to identify sentinel node metastases, but found on ultrasound-guided biopsy that many nodes were reactive hyperplasia and without metastatic disease [26]. They concluded that PET-CT is insufficient to diagnose regional metastases. Patients with advanced tumors should undergo workup for distant metastases with chest radiography, head CT, and abdominal ultrasound [27]. Enlarged, palpable nodes



were the primary indications in nine papers, high-grade tumors in two papers [28], and pathologic diagnosis of melanoma in one paper [26].

For all these reasons, the only really effective therapeutic weapon effective therapeutic weapon in the case of Marjolin's ulcer is preventive treatment, because it is the only one that allows to avoid the passage from a benign scar to a malignant and virulent tumor. This preventive treatment, which is based on the coverage of acute burns of deep acute burns, rehabilitation and the management of scars, and management of scars by compression. In order to avoid the development of hypertrophic and retractile sequelae that can lead to chronic ulceration, excision-grafting must be performed within two weeks of the deep burn following the deep burn in the "functional" areas, such as the joints, and functional" areas, such as joints, and within three weeks in other areas. As soon as healing is achieved, rehabilitation and compression are started. In the limbs, physical therapy performs active and passive mobilization in order to maintain joint amplitudes and prevent tissue retraction of the tissues. Compression involves the use of compression garments and the application of silicone plates, during the entire period of scar remodeling, 18 months to two years.

Massages performed by the physiotherapist, allow the scars to become more supple and to limit adhesions. Skin hydration with neutral emollients, prevents the skin from drying out, the appearance of cracks and the reopening of wounds.

## CONCLUSION

The occurrence of squamous cell carcinoma on burns scars is an infrequent but formidable complication. Its treatment is primarily preventive, based on correct initial management of burn lesions and regular monitoring of unstable scars. Moreover, in case of suspicious lesions, surgical treatment must be radical from the outset.

## REFERENCES

1. Hochberg, M., McNamara, K., & Kalmar, J. (2022). Marjolin's ulcer: A case report of a rare but preventable lesion. *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*, 133(5), e132.
2. Phillips, T. J., & Salman, S. M. (1998). Burn scar carcinoma. *Dermatol Surg*, 22, 561-565.
3. Gan, B. S., & Colcleugh, R. G. (1995). Melanoma arising in a chronic (Marjolin's) ulcer. *J Am Acad Dermatol*, 32, 1058-1059.
4. Ozek, C., Celik, N., Bilkay, U., Akalin, T., Erdem, O., & Cagdas, A. (2001). Marjolin's ulcer of the scalp: report of 5 cases and review of the literature. *The Journal of burn care & rehabilitation*, 22(1), 65-69.
5. Esther, R. J., Lamps, L., & Schwartz, H. S. (1999). Marjolin ulcers: secondary carcinomas in chronic wounds. *Journal of the Southern Orthopaedic Association*, 8(3), 181-187.
6. Kerr-Valentic, M. A., Samimi, K., Rohlen, B. H., Agarwal, J. P., & Rockwell, W. B. (2009). Marjolin's ulcer: modern analysis of an ancient problem. *Plastic and reconstructive surgery*, 123(1), 184-191.
7. Steffen, C. (1984). Marjolin's ulcer: Report of two cases and evidence that Marjolin did not describe cancer arising in scars of burns. *The American journal of dermatopathology*, 6(2), 187-194.
8. Hensel, K. S., Ono, C. M., & Doukas, W. C. (1999). Squamous cell carcinoma in chronic ulcerative lesions: a case report and literature review. *American Journal of Orthopedics (Belle Mead, NJ)*, 28(4), 253-256.
9. Nkere, U. U., & Banjar, A. (1997). Malignant change in a chronic skin lesion induced by an intravenous cannula. *Postgraduate medical journal*, 73(855), 45-47.
10. Berkwits, L., Yarkony, G. M., & Lewis, V. (1986). Marjolin's ulcer complicating a pressure ulcer: case report and literature review. *Archives of physical medicine and rehabilitation*, 67(11), 831-833.

11. Chlihi, A., Bouchta, A., Benbrahim, A., Bahechar, N., & Boukind, E. H. (2002, August). L'ulcère de Marjolin, destinée d'une cicatrice instable. À propos de 54 cas de séquelles de brûlure. In *Annales de chirurgie plastique esthetique* (Vol. 47, No. 4, pp. 291-297). Elsevier Masson.
12. Fleming, M. D., Hunt, J. L., Purdue, G. F., & Sandstad, J. (1990). Marjolin's ulcer: a review and reevaluation of a difficult problem. *The Journal of burn care & rehabilitation*, 11(5), 460-469.
13. Copcu, E., & Çulhaci, N. (2002). Marjolin's ulcer on the nose. *Burns*, 28(7), 701-704.
14. Kopp, J., Bach, A. D., Kneser, U., & Horch, R. E. (2006). Application of VAC-therapy during plastic surgical treatment of a bifocal Marjolin ulcer. *Zentralblatt für Chirurgie*, 131(Suppl. 1), S29-32.
15. García-Morales, I., Pérez-Gil, A., & Camacho, F. M. (2006). Marjolin's ulcer: burn scar carcinoma. *Actas Dermosifiliograficas*, 97(8), 529-532.
16. Simmons, M. A., Edwards, J. M., & Nigam, A. (2000). Marjolin's ulcer presenting in the neck. *The Journal of Laryngology & Otology*, 114(12), 980-982.
17. Hammond, J. S., Thomsen, S., & Ward, C. G. (1987). Scar carcinoma arising acutely in a skin graft donor site. *Journal of Trauma and Acute Care Surgery*, 27(6), 681-683.
18. Çeliköz, B., Demiriz, M., & Selmanpakoğlu, N. (1997). A shorter lag period of mesenchymal malignancy on Marjolin's ulcer. *Burns*, 23(1), 72-74.
19. Castillo, J. L., & Goldsmith, H. S. (1968). Burn scar malignancy in a possible depressed immune setting. *Surg Forum*, 19, 511-513.
20. Alcolado, J. C., Ray, K., Baxter, M., Edwards, C. W., & Dodson, P. M. (1993). Malignant change in dermatitis artefacta. *Postgraduate medical journal*, 69(814), 648-650.
21. Konigova, R., & Rychterova, V. (2000). Marjolin's ulcer. *Acta Chir Plast*, 42(3), 91-94.
22. Barr, L. H., & Menard, J. W. (1983). Marjolin's ulcer the LSU experience. *Cancer*, 52(1), 173-175.
23. Hill, B. B., Sloan, D. A., Lee, E. Y., McGrath, P. C., & Kenady, D. E. (1996). Marjolin's ulcer of the foot caused by nonburn trauma. *Southern medical journal*, 89(7), 707-710.
24. Eastman, A. L., Erdman, W. A., Lindberg, G. M., Hunt, J. L., Purdue, G. F., & Fleming, J. B. (2004). Sentinel lymph node biopsy identifies occult nodal metastases in patients with Marjolin's ulcer. *The Journal of burn care & rehabilitation*, 25(3), 241-245.
25. Shen, R., Zhang, J., Zhang, F., Du, Y., Liang, W., Xu, L., ... & Chen, X. (2015). Clinical characteristics and therapeutic analysis of 51 patients with Marjolin's ulcers. *Experimental and therapeutic medicine*, 10(4), 1364-1374.
26. Motamedolshariati, M., Rezaei, E., Beiraghi-Toosi, A., Jahani, A., Tayyebi Meibodi, N., Fattahi, A., & Sadeghi, R. (2015). Sentinel node mapping in Marjolin's ulcers: is it feasible. *Wounds*, 27(3), 54-62.
27. Tiftikcioglu, Y. O., Ozek, C., Bilkay, U., Uckan, A., & Akin, Y. (2010). Marjolin ulcers arising on extremities. *Annals of plastic surgery*, 64(3), 318-320.
28. Al-Zacko, S. M. (2013). Malignancy in chronic burn scar: A 20 year experience in Mosul-Iraq. *Burns*, 39(7), 1488-1491.